

# Are pre-MS stars older than we thought?

Tim Naylor & N.J. Mayne

School of Physics, University of Exeter, EX4 4QL, UK

**Abstract.** We present a consistent age ordering for young clusters and groups determined using the contraction of stars through their pre-main-sequence phase. We compare these with ages derived from the evolution of the upper main-sequence stars, and find the upper MS ages are older by a factor 1.5 to 2. We show that increasing the binary fraction and number of equal-mass binaries amongst the O-stars compared to the rest of the MS cannot remove this discrepancy.

**Keywords.** methods: statistical, stars: formation, open clusters and associations: general

In Mayne *et al.* (2007) and Mayne & Naylor (2008) we developed an age-ordering for young stars and groups based on the luminosity of the pre-main-sequence. Table 1 shows the resulting ages, including those derived in subsequent papers. In Naylor (2009) we derived ages by fitting the change position in the colour-magnitude diagram of upper-MS stars as they evolve from the zero-age MS to the terminal-age MS. We found that the MS ages are a factor 1.5 to 2 longer than the ages derived from the PMS.

After my presentation Gaspard Duchêne pointed out that binarism amongst O-stars is much higher than in the rest of the MS. A higher binary fraction will shift the centroid of the combined single-star and binary-star sequences redwards, mimicking an older age and perhaps explaining the older MS ages. As the mass-ratio distribution is equally important, we tested this idea using the most extreme assumption we could reasonably make, the strong hypothesis of Lucy (2006), which we approximated as 25% of binaries evenly distributed over  $0.95 < q < 1.0$ , and 75% evenly distributed over  $0.2 < q < 0.95$ . Using this, and a binary fraction (restricted to  $q > 0.2$ ) of 75% (e.g. Sana *et al.*, 2009) for all O-stars we find the ages of the clusters change by less than 5% compared with the results of Naylor (2009). Thus the discrepancy between the MS and PMS ages remains.

## References

- Jeffries, R.D., Naylor, T., Walter, F.M., Pozzo, M.P., & Devey, C.R. 2009, *MNRAS* 393, 538  
Lucy, L.B. 2006, *A&A* 457, 629  
Mayne, N.J. & Naylor, T. 2008, *MNRAS* 386, 261  
Mayne, N.J., Naylor, T., Littlefair, S.P., Saunders, E.S., & Jeffries, R.D. 2007, *MNRAS* 375, 1220  
Naylor, T. 2009, arXiv:0907.2307  
Naylor, T. & Jeffries, R.D. 2006, *MNRAS* 373, 1251  
Sana, H., Gosset, E., & Evans, C.J. 2009, arXiv:0909.0486

**Table 1.** Ages from PMS contraction

1Myr	2Myr	3Myr	4-5Myr	5-10Myr	10Myr	13Myr	40Myr
IC5146	ONC	$\lambda$ Ori,	IC348	$\gamma$ Vel <sup>2</sup>	NGC7160	h & $\chi$ Per	NGC2547
	NGC 6530	$\sigma$ Ori,	Cep OB3b <sup>1</sup>				
		NGC2264	NGC2362				

<sup>1</sup> Littlefair *et al.* in prep. <sup>2</sup> Jeffries *et al.* (2009)